

MANDELIN REAGENT INSTRUCTIONS

1. Carefully shake bottle before each use. Open the WIM Scientific Laboratories Mandelin Reagent's factory seal.
2. Using the provided mini tester spoon, place at least .010 to .005 Grams (**Tiny amount**) of the questionable substance into the empty testing vial.
3. Add one or two drops of the Mandelin Reagent into the testing vial. **The mandelin reagent is a strong yellow color.***
4. Watch carefully during the reaction time for color changes, any fizzing or smoking.
5. Refer to the **color chart (on back)** to determine what is present in the sample.
6. Rinse testing vial and the mini tester spoon thoroughly with soap and water after testing.
7. After successfully testing your substance, mini testing spoon and testing vial will need to be completely cleaned and dried before your next use.
8. After testing, the Mandelin Reagent bottle cap should be closed tightly and placed back into the bag to ensure no leakage or unwanted exposure occurs.
9. Also included are glow sticks and wristbands...because we love you. They may come in handy!



Mandelin Reagent Kits are **made to order** with manufacture dates stamped on the bag and will be useful for at least 3-6 months depending on proper storage. Keep out of direct sunlight and hot temperatures (Above 120 degrees) for best results and lasting usage.

Please note that a positive or negative reaction for any substance tested does not mean that a substance is safe. No chemical use is 100% safe. This will simply test for the presence of certain substances. A positive reaction for any compound still does not mean it's FULLY pure and could still be cut with other contaminants.

If reagent comes into contact with skin or eyes, wash promptly with soap and water. Latex gloves are highly recommended when handling kits.
If irritation persists, receive medical attention immediately. **KEEP OUT OF REACH OF CHILDREN.**



Only The Highest Quality Lab Grade Chemicals Used:
ISO 9001:2008 Quality Lab Material Certified




MANDELIN REAGENT REACTIONS

(Read all instructions on back first)*

SUBSTANCES:

Cocaine	Deep orange yellow
Ketamine	Deep reddish orange
MDMA/MDA	Dark bluish black
Methylone	Red
Amphetamine	Moderate bluish green
Methamphetamine	Dark yellowish green
Codeine	Dark olive
Heroin	Moderate reddish brown
Oxycodone	Darkish green
Mescaline	Dark yellowish brown
PMA/PMMA	Reddish brown
Salt (NaCl)	Strong orange



*Mandelin reagent is a very strong yellow color prior to being used for testing**



WARNING! PMA & PMMA ARE EXTREMELY DANGEROUS

WIM Scientific Laboratories LLC™

info@wimscilabs.com

